

Listening to the kids: the value of childhood palaeopathology for the study of rural Roman Britain

Article

Accepted Version

Rohnbogner, A. (2017) Listening to the kids: the value of childhood palaeopathology for the study of rural Roman Britain. *Britannia*, 48. pp. 221-252. ISSN 1753-5352 doi: <https://doi.org/10.1017/S0068113X17000149> Available at <http://centaur.reading.ac.uk/69645/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/S0068113X17000149>

Publisher: Society for the Promotion of Roman Studies

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Listening to the Kids: The Value of Childhood Palaeopathology for the Study of Rural Roman Britain

Anna Rohnbogner

ABSTRACT

Childhood palaeopathology remains an underutilised resource in the study of Roman Britain, particularly for exploring the lives of the rural population. Lesions in child skeletons provide unique insights into past lifeways and population health, as adverse environmental conditions translate more readily into the osteological record of these vulnerable members of society. To demonstrate the range of information gleaned from the children, 1,279 non-adults (0–17 years) from 26 first- to fifth-century urban and rural settlements were analysed, comparing morbidity and mortality in the most comprehensive study to date. The distribution of ages-at-death suggests migration between country and town, the latter presenting a stressful and unsanitary environment. However, as demonstrated by high rates of metabolic disease and infections, life in the countryside was hampered by demanding physical labour and potentially oppressive conditions with restricted access to resources.

Keywords: inequality deprivation; non-adult; migration; childhood pathology; resource stress; Roman rural settlement

INTRODUCTION: ROMANO-BRITISH CHILDHOOD

Human skeletal remains present the most direct form of evidence for studying past populations. Numerous palaeopathological studies of Romano-British adults have provided insight into population health, diet and lifeways, albeit with a bias towards urban cemeteries.¹ More recently, several studies have reported on health in both urban and rural contexts, yet the children did not receive emphasis.² Roman archaeology is no different from other branches of the discipline in its research concerns, therefore the childhood experience has not yet fully evolved as a dedicated subject in Roman scholarship. While stable isotope analysis of Romano-British children has received considerable attention,³ the actual health status of

children remains relatively unexplored and a comprehensive picture of growing up in Roman Britain is still lacking. As children are growing, they reflect adverse environmental conditions more readily than their parents.⁴ Child health is therefore a very powerful indicator of overall population health and dynamics, making non-adult (0–17 years) palaeopathology a primary source of evidence for past lifestyles and living environments. The osteological term ‘non-adult’ is not without problems but allows us to label, observe and discuss immature individuals in the past without involuntarily attaching social age norms.⁵ The cut-off age of 17 years is a product of the most accurate osteological ageing methods used and allows comparison between bioarchaeological datasets. Alongside age-at-death, a range of acquired conditions of ill-health can be observed in non-adult skeletons, which can provide information regarding cultural change, environmental pressures and life histories.

The classical literature on Romano-British children is limited and, as so often in the study of Romano-British lifeways, we are forced to look to Rome for analogies and guidance. The classical literature, epigraphic and iconographic references provide a wealth of information on the Roman childhood experience.⁶ However, particularly the written sources are based on individual accounts of a privileged minority, tainted with anecdotal undertones and very little to no mention of the working classes, peasantry and slaves.⁷ Inscriptions and imagery of Romano-British children are scarce and cannot fully communicate life histories of children from all orders of society within town and country.⁸ It is difficult to trace the everyday realities of a child in Rome, let alone at the north-western fringes of the Empire, post-dating most of the ancient literature by several centuries. Yet, awareness of the life course of Roman children as described in important texts, such as Soranus’ second-century A.D. *Gynaecology*, is beneficial.⁹ Soranus’ instructions on weaning practices,¹⁰ believed to have been followed in Rome, may be detectable in the osteological record of Romano-British children. However, in the absence of historical sources, direct skeletal evidence of ill-health in the children is the most comprehensive means of exploring the childhood experience in Roman Britain. This study aims to provide an overview of childhood pathology across the province, giving new insights into growing up in both town and country. Most importantly,

this analysis adds substantially to our current understanding of life in the understudied Romano-British countryside via the palaeopathology of the children.

CHILDREN FROM ROMAN ENGLAND: AN OVERVIEW OF HEALTH AND WELL-BEING

Acquired in childhood, cribra orbitalia and enamel hypoplasia have been addressed in skeletal populations across the Empire and Rome itself, allowing insight into general health, nutritional status and, more specifically, the prevalence of malaria.¹¹ High prevalence of malaria favours individuals with genetic mutations of resistance, which can manifest as β -thalassaemia and other anaemias.¹² Cribra orbitalia and porotic hyperostosis are frequently observed in Italian Roman populations, suggesting high rates of acquired and genetic anaemic conditions.¹³ Malaria is well recognised as an endemic disease in Classical Italy which impacted on morbidity and mortality to the extreme of yielding a demography of malaria, which may have also affected the southern areas of Roman Britain.¹⁴ In Italy, correlations between high rates of marrow hypoplasia and malarious areas have been identified. In Roman Britain however, cribrotic and enamel lesions as indicators of early childhood stress and a malarial threat were considerably less prevalent.¹⁵

Important insights into child health in Roman Dorset are provided in several palaeopathological studies which highlight a decline in child health from the Iron Age through to the late Roman period.¹⁶ Accounts of unprecedented detail of non-adult ill-health in Roman Britain stem from Lewis' work on metabolic disease, trauma, tuberculosis and thalassaemia from 364 children (0–17 years) from third- to fifth-century Poundbury Camp, Dorchester, Dorset.¹⁷ Elevated levels of metabolic disease at the site bear resemblance to those reported in post-medieval children.¹⁸ The Poundbury Camp cohort included three non-adults who were likely to have suffered from thalassaemia, providing direct osteological evidence for immigrants from the Mediterranean in Roman Dorchester.¹⁹ Particularly in the infants, high levels of blood-borne conditions and vitamin C and D deficiencies may be witness to early Christian fasting or ascetic practices adhered to by the women at the site.²⁰ We would also expect to see these lesions in children that suffered from thalassaemia major

and subsequently could not feed.²¹ Rickets was common at Poundbury Camp and has been interpreted as a result of swaddling, a Roman practice.²² However, the clinical literature demonstrates that swaddling is an overall beneficial childcare practice with a negligible effect on exposure to sunlight and therefore vitamin D deficiency.²³ Rather, a restricted diet for mothers and children, coupled with a weaning diet as described in Soranus²⁴ would have affected non-adult health at the site. The presence of childhood tuberculosis at Poundbury Camp indicates a tangible risk of infection in the children from Roman Dorchester.²⁵ Finding childhood tuberculosis confirms theories about urban living, such as crowded and unsanitary conditions, and suggests a depressed immune status of the children which allows for secondary infection and subsequent bony changes to occur.²⁶ As a singular site, Poundbury Camp exhibits an unexpectedly high degree of compromised child health.²⁷

Today, we believe that living in the countryside is beneficial in terms of both lifestyle and well-being and we tend to perceive this to be true of all time periods. However, rural life in Roman Britain will have come with significant health challenges due to the market economy and exploitative Imperial system.²⁸ Pitts and Griffin reported on compromised health in rural sites across central and southern England compared to urban cemeteries, characterised by higher levels of skeletal indicators of malnutrition, infection and strenuous physical activity.²⁹ Non-urban cemeteries exhibited fewer grave furnishings and greater inequality in the distribution of grave goods, possibly indicative of a relationship between living environment, social inequality and well-being in late Roman Britain. Redfern *et al.* studied urban-rural differences in first- to fifth-century burial grounds from Dorset.³⁰ The main findings included lower survivorship, higher mortality risk and smaller numbers of older adults in rural cemeteries. Excluding Poundbury Camp and associated cemeteries, individuals from rural sites showed higher rates of skeletal indicators of blood-borne disturbances and periosteal lesions which may relate to non-specific stress. These results challenge long-held perceptions of the benefits of rural living and highlight the need for a more dedicated study of health in the rural Romano-British population. We have to question what effect living and working arrangements had on the peasant population. Indeed, Romano-British society was highly stratified, where rich and poor would have collided in the

countryside with peasant farmers, or perhaps even bonded tenants, on one side and powerful land owners on the other. Were people ‘employed’ by benevolent villa owners, or would land management have been more similar to later Medieval manorialism, reliant on the exploitation of the workforce? This study seeks to demonstrate the important insights gleaned from the children and their palaeopathology, as a means to providing vital clues for discerning and conceptualising health in the countryside, social injustice and rural Romano-British lifeways.

MATERIALS AND METHODS

In the most comprehensive overview to date, 26 first- to fifth-century sites were included. The sample comprises 1,279 individuals (0–17 years), with 953 non-adult skeletons recorded by the author and osteological data from a further 316 gathered from (un)published reports (Table 1, FIG. 1). Three sites yielded a limited number of first-century burials (Winchester north, west and east, Springhead, Owslebury). Sources include the Museum of London Wellcome Osteological Database (WORD),³¹ published site reports and grey literature reports from the commercial sector. Poundbury Camp was deliberately excluded from analysis, since the site is not representative of non-adult health in major urban settlements as demonstrated in detail by Rohnbogner and Lewis.³² Sites were divided into major urban (*coloniae*, civitas capitals), minor urban (small towns, roadside settlements) and rural categories (*villae*, farmsteads, villages), following settlement classifications discussed in the literature.³³ Settlement classification in Roman Britain is complex and the terms major urban, minor urban and rural seek to acknowledge and describe the economic and administrative status of walled and unenclosed settlements, enabling comparisons in health across three distinct living environments. Age categories correspond with developmental stages and were used to minimise bias and allow cross-site comparison: perinate (<42 weeks gestation), 0.0–1.0 years, 1.1–2.5 years, 2.6–6.5 years, 6.6–10.5 years, 10.6–14.5 years and 14.6–17.0 years. The sample recorded by the author was aged using tooth crown and root formation stages,³⁴ long-bone length regression formulae³⁵ and skeletal fusion and maturation.³⁶ A variety of

pathological conditions described below were taken into consideration to explore ill-health in Romano-British children.

READING THE SIGNS – NON-ADULT PALAEOPATHOLOGY

A range of skeletal lesions may develop during childhood, as a result of blood-borne diseases, non-specific stress and infection, metabolic disease and trauma. The presence of pathological conditions, together with the age of the affected individual, allow observations on the prevalence of ill-health within a population. In turn, social, behavioural and biological factors can be explored, including child rearing practices, living environment and maternal health.

Blood-borne diseases

Pitting and porosity on the outer table of the skull (porotic hyperostosis) and orbital roof (cribra orbitalia), indicate blood-borne disorders. Non-adults display lesions more frequently than adults due to a reduced capacity of sustaining higher red blood cell production.³⁷ The palaeopathological literature has ascribed cribra orbitalia and porotic hyperostosis to iron deficiency anaemia, arising from excessive blood loss, insufficient dietary iron intake, intestinal parasites or diarrhoea.³⁸ Lesions also stem from vitamin B12/B9 deficiency in megaloblastic anaemias which may co-occur with iron deficiency.³⁹ The informative value of cribra orbitalia and porotic hyperostosis is considerable, including general childhood health, maternal health and feeding practices. A largely plant-based diet will leave the mother deficient in vitamin B12 and iron. In turn, the breastfed infant is nursed on milk deficient in vitamins and minerals. Prolonged breastfeeding without adequate supplementation in the growing child will eventually lead to deficiency. Lastly, haemolytic anaemias, such as thalassaemia, are recognisable in the osteological record.⁴⁰

Stress and infection

Enamel hypoplasia

Enamel hypoplasia are linear bands or pits in tooth crown enamel, a retrospective record of early childhood stress.⁴¹ Defects can affect all teeth and are most often recorded on the crowns of the anterior permanent dentition, formed until the age of four.⁴² Generally, enamel hypoplasia are a non-specific indicator of poor health and stressors may include trauma, mal- or undernutrition, fever and infection.⁴³ Deciduous canines may be affected by unilateral pitted enamel defects, arising from weakened bone structure coupled with minor trauma from normal motor development during infancy. Their aetiology is complex and probably not linked with a systemic cause.⁴⁴

Non-specific infection

Endocranial lesions are found on the inner table of the skull.⁴⁵ Lesions may occur secondary to infection, trauma, tumours or meningitis and have been observed in cases of tuberculosis and vitamin deficiencies.⁴⁶ Sub-periosteal new bone formation is recognisable as raised, porous patches of new bone.⁴⁷ This type of lesion is an inflammatory response to infection, trauma, circulatory disorders, joint disease, haematological disease, skeletal dysplasia and metabolic or neoplastic disease.⁴⁸

Tuberculosis

Tuberculosis is a disease of malnutrition, poverty and overcrowding. It is a chronic infectious disease of the lungs, skin, lymph nodes, intestines and, in rare cases, bones and joints.⁴⁹ The bacterial genus specific to humans, *Mycobacterium tuberculosis*, is spread via airborne droplets, sputum or human waste and can be transmitted from mother to foetus.⁵⁰ Ingesting infected animal products will spread the animal equivalent bacillus, *Mycobacterium bovis*, from bovines to humans.⁵¹

Metabolic disease

Vitamin D deficiency

The prohormone vitamin D is absorbed via the intestine or formed by the skin's dermal cells in response to ultraviolet light and is vital to the formation and maintenance of healthy bone

structure.⁵² In children, especially infants, chronic vitamin D deficiency has its most marked impact, leading to rickets and osteomalacia. Rachitic children exhibit unmineralised bone that is porous in appearance and, when mechanical forces are applied, is prone to characteristic bending deformities. Apart from exposure to sunlight and associated cultural practices, rickets also informs on calcium deficiency, female health and patterns in transitional feeding.⁵³

Scurvy

The clinical condition of scurvy results from chronic vitamin C deficiency. Non-adult skeletons, especially those of infants and young children are more likely to exhibit scorbutic lesions due to rapid growth.⁵⁴ Skeletal growth is slowed down and sub-periosteal haematomas occur at weakened walls of small blood vessels, yielding characteristic porous patches of new bone, particularly in the skull.⁵⁵ Apart from direct evidence for a lack of fresh fruits and vegetables, scurvy in archaeological populations is a vehicle for exploring resource stress, social hierarchies, ecology and behaviour.⁵⁶

Trauma

Dislocation, alteration to the shape of the bone, partial and complete breaks are distinct skeletal injuries observable in the osteological record.⁵⁷ Childhood fractures are a vital tool for providing information on occupational stress and exposure to trauma. Paediatric bone may yield distinctive fractures that can quickly lose visibility in the palaeopathological record due to fast remodelling.⁵⁸

The following methods were used by the author. Crude prevalence of enamel hypoplasia were recorded macroscopically.⁵⁹ Infections of a non-specific nature were recorded using sub-periosteal new bone formation and endocranial lesions.⁶⁰ Tuberculosis was diagnosed in the presence of lytic lesions in the spine with visceral rib lesions and widespread new bone formation.⁶¹ Cribra orbitalia and porotic hyperostosis were scored according to Stuart-Macadam.⁶² A diagnosis for thalassaemia was based on the presence of

costal osteomas or 'rib-within-a-rib',⁶³ osteopenia and generalised cortical thickening, cribra orbitalia and porotic hyperostosis.⁶⁴ Vitamin D deficiency was identified following the criteria of Ortner and Mays, Ortner, and Brickley and Ives.⁶⁵ Vitamin C deficiency was scored according to Brickley and Ives⁶⁶ and the 'Ortner criteria' for lesions on the skull and metaphyseal ends of long bones.⁶⁷

As expected, working with secondary palaeopathological data presents limitations, as recording methods and reporting standards may differ. The Chi-squared 3x2 non-parametric statistic (X^2) was used to test for differences between groups. False positives were avoided by setting the confidence interval at 99.5 per cent ($p=0.005$) and using the test sparingly only when percentages indicated great differences between groups.

RESULTS

Age-at-death

There were significantly fewer perinates (16.9 per cent, $n=88$; $X^2=21.71$, d.f.=2, $p<0.001$) and infants (15.9 per cent, $n=83$; $X^2=38.25$, d.f.=2, $p<0.001$) reported from major urban sites. Most striking were the high numbers of rural perinates (29.9 per cent, $n=95$) and infants from minor urban sites (32.9 per cent, $n=145$). In the 6.6–10.5 year old group, significantly more individuals come from major urban cemeteries than elsewhere (12.6 per cent, $n=66$; $X^2=14.98$, d.f.=2, $p<0.001$). The 14.6–17.0 year olds were significantly more frequent in major urban cemeteries (10.3 per cent, $n=54$; $X^2=20.25$, d.f.=2, $p<0.001$) compared with minor urban sites (2.9 per cent, $n=13$) (Table 2, FIG. 2).

The pathology

The distribution of lesions across the settlement types was not uniform (Table 3). Elevated rates of cribra orbitalia and porotic hyperostosis, thalassaemia, enamel hypoplasia and tuberculosis were found in the major urban cohort (FIG. 3). No additional cases of probable thalassaemia were reported, other than those already discussed in the literature.⁶⁸ Infants were excluded from analysis for infectious and metabolic diseases, as their bones are increasingly reactive and porous, mimicking lesions of pathological origin.⁶⁹ Deficiency diseases of

vitamin D and C in children aged 1.1 years and older were most frequent in the rural sample at 6.9 per cent (n=10) (FIG. 3).

Haematopoietic lesions

High rates of cribra orbitalia and porotic hyperostosis were observed in the rural children aged 1.1–2.5 (29.4 per cent, n=10) and 6.6–10.5 years old (27.8 per cent, n=5). The rates reported from minor urban sites were elevated in 2.6–6.5 year olds (27.1 per cent, n=19), where rates from major urban and rural sites were similar at 18.8 per cent (n=16) and 16.3 per cent (n=7) respectively (Table 4, FIG. 4).

Stress and infection

Enamel defects were significantly more frequent in major urban sites (15.9 per cent, n=83; $\chi^2=29.43$, d.f.=2, $p<0.001$). Enamel hypoplasia in deciduous canines were reported in urban non-adults up to 2.5 years old. In 6.6–10.5 year olds, rates are similar in the major urban (27.3 per cent, n=18) and rural sites (27.8 per cent, n=5). From 10.6 years and older, the rural enamel hypoplasia rate is lower than in the urban sites, although not statistically significant (Table 5, FIG. 5). Endocranial lesions and sub-periosteal new bone formation occurred at similar rates in major urban (15.1 per cent, n=53) and rural contexts (14.5 per cent, n=21), and were most prevalent in the minor urban sample (20.4 per cent, n=43). Children from minor urban contexts showed the highest rate amongst the 1.1–2.5 year olds (26.3 per cent, n=15). In 2.6–6.5 and 6.6–10.5 year olds, lesions occurred at similar rates in the minor urban and rural sites at rates of 20.0 per cent (n=14)/18.6 per cent (n=8) and 22.6 per cent (n=7)/22.2 per cent (n=4). In 10.6–14.5 and 14.6–17.0 year olds, major urban rates were highest. In 14.6–17.0 year olds, the rural rate was low at 4.5 per cent (n=1), although the result was not statistically significant (Table 6, FIG. 6).

Tuberculosis

Individuals with lesions that are either strongly suggestive or possible tuberculosis are few (n=9). Suspected and probable cases were reported from all three site types, with a bias

towards major urban settlements (0.8 per cent, n=4).⁷⁰ A further two probable cases were identified at minor urban Ancaster and Ashton and one possible case at rural Cannington.⁷¹

Metabolic disease

Vitamin D deficiency was reported in 14 individuals (2.0 per cent) of all ages, except amongst 10.6–14.5 year olds. Rickets was most frequent in the 1.1–2.5 year age group (5.5 per cent, n=8), where it was most prevalent in major urban sites (7.3 per cent, n=4) and lowest in minor urban settlements (3.5 per cent, n=2). In the rural sample, vitamin D deficiency was only reported in 1.1–2.5 year olds at 5.9 per cent (n=2). An adolescent female from Lankhills, Winchester, was the oldest individual with vitamin D deficiency. Lesions were healed implying that rickets occurred earlier in childhood (Table 7, FIG. 7).⁷²

A total of 16 individuals (2.3 per cent) across all ages were reported as vitamin C deficient. Prevalence was highest in 1.1–2.5 year olds (3.4 per cent, n=5), where 8.8 per cent (n=3) of rural individuals were affected and only 1.8 per cent (n=1) of non-adults from major urban and minor urban settlements. The distribution was not significant ($X^2=3.84$, d.f.=2). In the minor urban sample, scurvy was absent from the age of 2.6–6.5 years. The rural cohort exhibited higher prevalence rates across all age groups. The oldest affected individual was a 14.6–17.0 year old from Bath Gate, Cirencester, with healed lesions (Table 7, FIG. 8).

Trauma

Trauma affected 17 individuals from all three settlement types, the highest rate at 1.6 per cent (n=5) in the rural sample. The majority of trauma was identified in children aged 10.6 years and older (64.7 per cent, n=11). Higher fracture rates were seen in 10.6–14.5 year olds from minor urban sites. Rural 14.6–17.0 year olds exhibited the highest fracture rate at 9.1 per cent (n=2) (Table 8). The left side (n=9) was more often affected than the right (n=5). Clavicular fractures were most frequent (29.4 per cent, n=5), followed by tibial (17.6 per cent, n=3) and radial fractures (11.8 per cent, n=2), the latter most common in the major urban sample (n=2, 28.6 per cent).⁷³ Minor urban non-adults exhibited similar fracture rates of the clavicle, long

bones of the lower limbs and hand/foot phalanges (FIG. 9). Clavicular fractures were most frequent in rural non-adults (60.0 per cent, n=3) (Table 9).

Falls on to the shoulder, outstretched arm or hand may cause fractures to the clavicle, humerus, radius or ulna (FIGS 10 and 11).⁷⁴ Tibial and femoral fractures require considerable force either through violent injury, a direct blow to the element or a strong twisting force.⁷⁵ The youngest individual with trauma was a Winchester infant with a healing rib fracture (FIG. 12). The fracture may have arisen due to direct trauma,⁷⁶ complications in labour,⁷⁷ be a stress fracture from repeated coughing or vomiting,⁷⁸ or even be caused by accidental, over-zealous swaddling and infant handling.⁷⁹

DISCUSSION

This study is the largest scale analysis of non-adult health in Roman Britain to date, specifically focusing on morbidity and mortality of 0–17 year olds from urban and rural contexts. Achieving as broad an overview as possible, the study relies on data compiled through primary analysis by the author or on relevant data drawn from (un)published reports and databases (secondary data). This approach presented challenges. The secondary data is flawed due to inconsistent methods and variation in recording. Hence, the study provides a broad-brush approach, yielding observations of initial trends that can be explored in more detail using the primary data alone.⁸⁰ Osteological analysis is reliant on individuals that have died and were buried at a particular site⁸¹ and can never be truly representative of all individuals that have lived and died in major urban, minor urban or rural settlements. The chronology of the sample means that the majority of sites date to the later phases of Roman Britain when inhumation was the more common funerary rite. Limitations are also introduced by excavation and preservation bias, dictating the spatial and geographical spread of the sample. The north is poorly represented with only one major urban site, Trentholme Drive, York. The West Country is represented by a cluster of rural sites in Dorset and Somerset. Instead, the focus is primarily on the central and, to a lesser extent, southern areas of Roman Britain (FIG. 1). The rural sample is comparatively small, a product of fewer commercial investigations of rural sites, a focus on urban archaeology and the possibility of a bias

towards cremation rites on rural sites. In the adolescents aged 14.6–17.0 years, we have to consider the possibility of an artificial peak due to different ageing techniques followed in reports and the subsequent inclusion of older individuals. Some interesting trends in demography, stress and metabolic disease between the urban and rural groups have become apparent. Age-at-death and the overall distribution of enamel hypoplasia reached statistical significance between urban and rural groups at 99.5 per cent confidence while a peak in vitamin C deficiency is apparent in the rural children.

Infant mortality – reality or mirage?

High levels of perinatal burials in the rural sample and high representation of infants in minor urban cemeteries were witnessed. Neonatal deaths are caused by endogenous (i.e. biological and genetic) factors and reflect problems inherent at birth, which are most commonly seen in rural groups today.⁸² Scott and Duncan suggested that environmental factors, including infection, nutrition, poisoning and accidents are decisive in post-neonatal mortality.⁸³ These would reflect different stresses experienced by mothers and babies in urban and rural environments in Roman Britain. Previous studies have shown that urbanisation and increased industrialisation in past populations triggered elevated post-neonatal mortality.⁸⁴ The overcrowding, pollution and infection seen in major urban environments would have presented greater risks for post-neonatal mortality.⁸⁵ The results for perinatal and infant mortality suggest that these dangers did not exist in Romano-British urban sites. This seems unlikely and differences are probably due to differential infant burial rites and recovery.⁸⁶ Some rural sites have high proportions of neonate burials, often recovered from within the settlement boundaries, including Owslebury, Hampshire (43.8 per cent of total non-adult burials), Catsgore, Somerset (52.6 per cent), and Frocester (81.4 per cent) and Huntsman's Quarry (83.3 per cent) in Gloucestershire. The large percentage of perinates reported at Frocester is even more remarkable for the total absence of older infants, suggesting this may be a designated perinatal burial site. Some major urban cemeteries had low numbers of perinates, particularly the London burial grounds (4.0 per cent) and Butt Road, Colchester (2.8 per cent), with no perinates at Trentholme Drive, York, and the Gloucester cemeteries.

Low representation of infants in managed cemeteries is not exclusive to Roman Britain.⁸⁷ Distinct rites for those dying within the first year of life may have been a contemporary ritual practised across pockets of the Empire. Pliny the Elder describes children as lacking a soul until the age of teething at around six months old,⁸⁸ possibly accounting for differential treatment in death.⁸⁹ Funerary practice may have dictated the interment of babies and infants within the settlement boundaries, in clusters in a dedicated area of the cemetery, or at a separate site altogether, perhaps even in ‘infant corners’ that are yet to be excavated, ultimately impacting on the urban and rural rates of infant mortality observed in this study.⁹⁰

The weanling’s dilemma

Pathological lesions in children up to 2.5 years old attest to inadequate strategies in supplementary feeding. Insufficient calcium in urban weanlings may have been a problem, as supported by the highest rate of rickets in 1.1–2.5 year olds from major urban sites (7.3 per cent). Calcium deficiency prompting rickets is usually seen after the age of two years old following inadequate weaning strategies.⁹¹ However, as we cannot differentiate between rickets caused by either calcium or vitamin D deficiency, we have to consider the effect of clothing, childcare practices and living environment.⁹² Mothers in major urban towns may have kept their children indoors to recover and rest. Subsequently, vitamin D deficiency ensued, possibly exacerbated by low calcium levels in the diet, prolonged breastfeeding and gastrointestinal maladies that limited calcium absorption and ultimately caused rachitic lesions.⁹³

A slight increase in mortality is observed in 1.1–2.5 year olds from minor urban sites, which may correspond with the peak in infection observed in this age group and attest inadequate weaning strategies.⁹⁴ Rural weanlings demonstrate elevated levels of scurvy, cribra orbitalia and porotic hyperostosis when compared to their urban counterparts. Suitable foods may have been scarce. Vitamin C deficiency may have been a product of cereal-based supplementary foods, cooking/boiling of fresh produce and nursing by a deficient mother or wet nurse.⁹⁵ Anaemia may have been prompted by infection, high pathogen load and parasites introduced by a diet low in iron and unhygienic feeding practices.⁹⁶

Implications of childhood tuberculosis

Generally, evidence for infection was present in all three site types. Reporting of tuberculosis was low overall, but reinforces notions of a crowded and unsanitary urban environment.⁹⁷ Low-quality air may have been a problem in both town and country, as demonstrated by skeletal lesions indicative of respiratory infections such as bronchitis.⁹⁸ Smoke off the hearths and poor ventilation would have caused respiratory disorders in rural dwellings, while close contact with livestock would have caused transmission of *T. bovis* to humans.⁹⁹ The effects of childhood tuberculosis in Roman Britain may have been felt widely. Blindness, deafness and mental retardation follow when the initial infection spreads to the meninges of young children under four years old.¹⁰⁰

Adolescence and working lives

Proportionately higher rates of 6.6–10.5 and 14.6–17.0 year olds were observed in major urban settlements. Their lower numbers in non-urban cemeteries indicates that older children may have migrated to major urban settlements. Infections of non-specific origin were on the rise in rural children until the age of 6.6–10.5 years, after which they declined from 22.2 per cent to 4.5 per cent in 14.6–17.0 year olds. An inverse pattern was apparent in the major urban cohort, with infections among the lowest until 6.6–10.5 years, followed by an increase to 18.5 per cent in 14.6–17.0 year olds. This opposite relationship further supports the hypothesis of the migration of older children into the towns. Migration between rural and urban populations has been suggested previously in regional studies from Dorset and London.¹⁰¹ The urban environment exposed incoming children to new pathogens and demanding physical labour, eventually raising mortality.¹⁰² Although there is limited documentary evidence, the existence of child (forced) labour and a young start to the working life is feasible.¹⁰³ Children from as young as five years old would have been tasked with household chores, with working life commencing in the early teens or even pre-teens.¹⁰⁴

The majority of the rural population would have performed strenuous agricultural labour.¹⁰⁵ Workers needed adequate nutrition for energy expenditure in this physically

demanding job, while also being exposed to accidents and infectious diseases via livestock. Fracture locations in the rural cohort may bear witness to working with traction animals. Spinal degeneration in a 16-year-old female from Cannington further demonstrates the extent of intense physical activity.¹⁰⁶ Although enamel hypoplasia affected fewer individuals in the rural cohort, rates soared in the 6.6–10.5 year age group to the highest in the sample (27.8 per cent). Rural children who were strong enough to survive adverse conditions in early childhood faced an event aged 6.6–10.5 years old that caused elevated mortality. Non-specific infection, cribra orbitalia and porotic hyperostosis were all higher in the rural 6.6–10.5 year olds, compared to their major urban peers. We may suggest that the higher incidence of ill-health in this age group marks the biological signature of a social age transition whereby children commenced their working life. This tallies with what is known from historical sources regarding the shifting social status of children from the age of around seven years.¹⁰⁷

Apart from agricultural labour, industrial activity such as quarrying may have taken place on some rural sites, including Cannington, Somerset,¹⁰⁸ possibly accounting for the rise in mortality in 14.6–17.0 year olds from rural cohorts (6.9 per cent). Equally, the major urban towns were constantly under construction, with some buildings unsafe.¹⁰⁹ Indeed, five of the seven (71.4 per cent) fractures sustained by non-adults from major urban sites may stem from falls. Additionally, the densely populated poor districts and close proximity of living and working space would have exposed adolescent inhabitants to health risks in large towns.¹¹⁰

Diseases of deprivation?

Unexpectedly, rickets and scurvy were present in rural children. Since rickets was only found in 1.1–2.5 year olds, it is linked to calcium deficiency in young rural children or childcare practices. Perhaps these included shielding the poorly child from the sun. Scurvy was found in all but the oldest age group in the rural sample. Vitamin C levels in the body are affected by reduced nutritional intake, increased requirements, malabsorption or genetic causes.¹¹¹ Especially in young children, fussy eaters, religious dietary practices, low socioeconomic background, neglect, infection, inflammation, anaemia, gastrointestinal diseases, infections

and deficiencies in the pregnant and breastfeeding mother have to be considered.¹¹² Scorbutic lesions not only indicate absence of certain foods, but also more widespread under- or malnutrition secondary to food insecurity, preferential feeding and shortcomings in subsistence economy.¹¹³ Rural children suffered with haematopoietic stress, demonstrated by high rates of cribra orbitalia and porotic hyperostosis in 1.1–2.5 and 6.6–10.5 year olds. Children would have sustained chronic health insults in rural environments, with nutritional stress, infections and pathogens and parasites affecting both children and their mothers.¹¹⁴

The literature on dietary variation in Roman Britain indicates that the consumption of animal and plant foods was not uniform across the social strata. Meat was especially less accessible to lower-status individuals.¹¹⁵ A cereal-based diet introduces excessive amounts of phytates which inhibit intestinal absorption of iron.¹¹⁶ The haematopoietic and metabolic deficiencies observed in the rural children may therefore attest to lower status. Vitamin C deficiency is prevented by ingesting small amounts of fruits and vegetables, such as a cup of leek or pear each day.¹¹⁷ Social change and status differences dictate the foods people have access to, affecting those at the bottom of the social ladder most profoundly.¹¹⁸ Populations experience a negative impact on health during rapid economic change or modernisation.¹¹⁹ For Roman Britain, or England more specifically, malnutrition in the rural children may be a result of landownership, affecting food security and distribution.¹²⁰ The Roman conquest was followed by a rapid population increase and extensive reorganisation of the rural landscape.¹²¹ The rural population was forced to adapt, supply for the urban population and army and provide produce for trade and taxes.¹²² Agricultural production then further increased towards the end of the Roman rule.¹²³ During the fourth century, taxation changed from money into kind, tenants were legally tied to the estate and land tenancy became hereditary.¹²⁴ The rural population was under increasing strain, while landowners were relieved of having to provide food and accommodation to their (bonded) workforce.¹²⁵ Patronage systems may be seen from the fourth and fifth centuries, in some respects similar to later feudalism in Medieval Europe.¹²⁶ Pressure mounted due to events on the continent, including famine in the Rhineland.¹²⁷ Britain had become one of the main exporters of crops to the Rhine army, which further increased demand for produce while manpower remained

the same.¹²⁸ Additionally, the church as a powerful new landowner has to be considered, negotiating land distribution, tenancy and taxation.¹²⁹ Ultimately the rural population felt the strain, as the part of society from which resources are extracted is most affected by political instability, economic change and food shortages.

CONCLUSION

This study has presented palaeopathological and age-at-death data for 1,279 non-adults from 26 Romano-British sites, providing the most comprehensive survey of children from urban and rural Roman Britain to date. The data allow us to deliberately contrast ill-health between town and country, albeit with a bias towards later Roman phases and sites from central and southern England. This provides promising new insights into Romano-British life, from the possibility of distinct burial rites for the death of the youngest, to the movement of children from rural to urban areas and the start of their working lives. The study also highlights the adversity faced by some children, the impact on their health and, most importantly, some distinct differences in the stressors between urban and rural environments. Rural settlements posed threats to the well-being of children, certainly at a more pressing scale than expected, and with important implications for our current understanding of everyday lived realities in rural Roman Britain. To date, the vast majority of Roman Britons living in the countryside have remained archaeologically silent, even more so the women and children, similar to the ‘voiceless poor’¹³⁰ in developing and developed countries today. The findings presented in this article not only suggest a certain kind of lifestyle for the rural population, but also urge us to reconsider current ideas of living and working in Roman Britain and how we source our understanding of daily life for those outside the urban centres. Social stratification of Romano-British society may be to blame and should remain on the research agenda to gauge its full effect on population health and well-being.

ACKNOWLEDGEMENTS

I am particularly grateful to Drs Mary Lewis and Hella Eckardt for their interest, suggestions and comments. This research would not have been possible without access to skeletal

collections and I am indebted to Robert Kruszynski, Natural History Museum London; Keith Fitzpatrick-Matthews, North Hertfordshire District Council Museums Service; Dr. Simon Mays, English Heritage/Historic England; Christiane Jeuckens and David Moon, Oxfordshire Museums Service; Dr. Jo Buckberry, Biological Anthropology Research Centre at the University of Bradford; Steve Minnitt, Somerset Heritage Centre; David Allen and Helen Rees, Hampshire Cultural Trust; Sarah Wilson, Peterborough Museum; Lorraine Cornwell, Rutland County Museum; Dr. Alison Brooks and Heather Dawson, Corinium Museum; Dr. Paul Sealey and Jessica Dowdell, Colchester and Ipswich Museums; Timothy Vickers, Luton Culture. FIG. 9 was adapted and reproduced with kind permission from Dr Mirjana Roksandic. I want to thank David Moon at Oxfordshire Museums Service, Jessica Dowdell at Colchester and Ipswich Museum, and Helen Rees at Hampshire Cultural Trust, for granting permission for photography and publication of FIGS 10, 11 and 12. This research was funded by an AHRC PhD studentship.

Department of Archaeology, University of Reading

a.j.rohnbogner@reading.ac.uk

BIBLIOGRAPHY

- Armelagos, G.J., Sirak, K., Werkema, T., and Turner, B.L. 2014: 'Analysis of nutritional disease in prehistory: the search for scurvy in Antiquity and today', *International Journal of Palaeopathology* 5, 9–17
- Ayi, K., Turrini, F., Piga, A., and Arese, P. 2004: 'Enhanced phagocytosis of ring-parasitized mutant erythrocytes: a common mechanism that may explain protection against falciparum malaria in sickle trait and beta-thalassemia trait', *Blood* 104, 3364–71
- Barnett, C., McKinley, J.I., Stafford, E., Grimm, J.M., and Stevens, C.J. 2011: *Settling the Ebbsfleet Valley: High Speed 1 Excavations at Springhead and Northfleet, Kent. The Late Iron Age, Roman, Saxon and Medieval Landscape. Volume 3: Late Iron Age to Roman Human Remains and Environmental Reports*, Oxford/Salisbury

- Barsness, K.A., Cha, E., Bensard, D., Calkins, C.M., Partrick, D.A., Karrer, F.M., and Strain, J.D. 2003: 'The positive predictive value of rib fractures as an indicator of nonaccidental trauma in children', *The Journal of Trauma Injury, Infection, and Critical Care* 54, 1107–10
- Booth, P., Simmonds, A., Boyle, A., Clough, S., Cool, H.E.M., and Poore, D. 2010: *The Late Roman Cemetery at Lankhills, Winchester. Excavations 2000–2005*, Oxford
- Boyle, A., and Early, R. 1999: *Excavations at Springhead Roman Town, Southfleet, Kent*, Oxford Archaeological Unit Occasional Paper 1, Oxford
- Bradley, K.R. 1984: *Slaves and Masters in the Roman Empire: A Study in Social Control*, Oxford
- Bradley, K.R. 1991: *Discovering the Roman Family: Studies in Roman Social History*, Oxford
- Breeze, D.J. (ed.) 2014: *The Impact of Rome on the British Countryside. A Conference Organised by the Royal Archaeological Institute, Chester, 11–13 October 2013*, London
- Brickley, M., and Ives, R. 2006: 'Skeletal manifestations of infantile scurvy', *American Journal of Physical Anthropology* 129, 163–72
- Brickley, M., and Ives, R. 2008: *The Bioarchaeology of Metabolic Bone Disease*, London
- Brothwell, D.R., Powers, R., and Hirst, S.M. 2000: 'The pathology', in P. Rahtz, S. Hirst and S.M. Wright. *Cannington Cemetery: Excavations 1962–3 of Prehistoric, Roman, Post-Roman, and Later Features at Cannington Park Quarry, near Bridgwater, Somerset*, Britannia Monograph Series 17, London, 195–239
- Buckley, H.R., Kinaston, R., Halcrow, S.E., Foster, A., Spriggs, M., and Bedford, S. 2014: 'Scurvy in a tropical paradise? Evaluating the possibility of infant and adult vitamin C deficiency in the Lapita skeletal sample of Teouma, Vanuatu, Pacific Islands', *International Journal of Paleopathology* 5, 72–85
- Bulloch, B., Schubert, C.J., Brophy, P.D., Johnson, N., Reed, M.H., and Shapiro, J.A. 2000: 'Cause and clinical characteristics of rib fractures in infants', *Pediatrics* 105, e48

- Burleigh, G.R., and Fitzpatrick-Matthews, K.J. 2010: *Excavations at Baldock, Hertfordshire, 1978–1994. Volume 1: An Iron Age and Romano-British Cemetery at Wallington Road, Letchworth*
- Burn, A.R., 1970: *The Romans in Britain: An Anthology of Inscriptions*, Oxford
- Carroll, M., and Graham, E. (eds) 2014: *Infant Health and Death in Roman Italy and Beyond*, Journal of Roman Archaeology Supplementary Series 96, Portsmouth, RI
- Chambers, R.A. 1987: 'The late and sub-Roman cemetery at Queenford Farm, Dorchester-on-Thames, Oxon', *Oxoniensia* 52, 35–70
- Clough, S., and Boyle, A. 2010: 'Inhumations and disarticulated human bone' in Booth *et al.* 2010, 339–99
- Collis, J.R. 1968: 'Excavations at Owslebury, Hampshire: an interim report', *Antiquaries Journal* 48, 18–31
- Collis, J.R. 1977: 'Owslebury (Hants) and the problem of burials on rural settlements', in R. Reece (ed.), *Burial in the Roman World*, Council for British Archaeology Research Report 22, London, 26–34
- Cool, H.E.M. 2006: *Eating and Drinking in Roman Britain*, Cambridge
- Crandall, J.J. 2014: 'Scurvy in the Greater American Southwest: modelling micronutrition and biosocial processes in contexts of resource stress', *International Journal of Palaeopathology* 5, 46–54
- Crandall, J.J., and Klaus, H.D. 2014: 'Advancements, challenges and prospects in the paleopathology of scurvy: current perspectives on vitamin C deficiency in human skeletal remains', *International Journal of Palaeopathology* 5, 1–8
- Crummy, N. 2010: 'Bears and coins: the iconography of protection in late Roman infant burials', *Britannia* 41, 37–93
- Crummy, N., and Crossan, C. 1993: 'Excavations at Butt Road Cemetery, 1976–9, 1986 and 1989', in N. Crummy, P. Crummy and C. Crossan, *Excavations of Roman and Later Cemeteries, Churches and Monastic Sites in Colchester, 1971–88*, Colchester Archaeological Report 9, Colchester, 4–163

- Cummings, C. 2009: 'Meat consumption in Roman Britain: the evidence from stable isotopes', in M. Driessen, S. Heeren, J. Hendriks, F. Kemmers and R. Visser (eds), *TRAC 2008: Proceedings of the Eighteenth Annual Theoretical Roman Archaeology Conference*, Oxford, 73–83
- Dix, B. 1983: 'Ashton Roman town', *South Midlands Archaeology* 13, 18–20
- Djuric, M., Milovanovic, P., Janovic, A., Draskovic, M., Djukic, K., and Milenkovic, P. 2008: 'Porotic lesions in immature skeletons from Stara Torina, Late Medieval Serbia', *International Journal of Osteoarchaeology* 18, 458–75
- Durham, B., and Rowley, T. 1972: 'A cemetery site at Queensford Mill, Dorchester', *Oxoniensia* 37, 32–8
- Esmonde Cleary, S. 2000: 'Putting the dead in their place: burial location in Roman Britain', in J. Pearce, M. Millett and M. Struck M (eds), *Burial, Society and Context in the Roman World*, Oxford, 127–149
- Esmonde Cleary, S. 2004: 'Britain in the fourth century', in Todd 2004, 409–27
- Facchini, F., Rastelli, E., and Brsili, P. 2004: 'Cribra orbitalia and cribra cranii in Roman skeletal remains from the Ravenna area and Rimini (I–IV century AD)', *International Journal of Osteoarchaeology* 14, 126–36
- Faith, R. 1997: *The English Peasantry and the Growth of Lordship*, London
- Farmer, P. 2008: 'Challenging orthodoxies: the road ahead for health and human rights', *Health and Human Rights* 10, 5–19
- Fewtrell, M.S., Morgan, J.B., Duggan, C., Gunnlaugsson, G., Hibberd, P.L., Lucas, A., and Kleinman, R.E. 2007: 'Optimal duration of exclusive breastfeeding: what is the evidence to support current recommendations?' *American Journal of Clinical Nutrition* 85, 635–8
- Foote, K., and Marriott, L. 2003: 'Weaning of infants', *Archives of Diseases in Childhood* 88, 488–92
- Frenzen, P.D., and Hogan, D.P. 1982: 'The impact of class, education, and health care on infant mortality in a developing society: the case of rural Thailand', *Demography* 19, 391–408
- Fulford, M. 2004: 'Economic structures', in Todd 2004, 309–26.

- Fuller, B.T., Molleson, T.I., Harris, D.A., Gilmour, L.T., and Hedges, R.E.M. 2006: 'Isotopic evidence for breastfeeding and possible adult dietary differences from Late/Sub-Roman Britain', *American Journal of Physical Anthropology* 129, 45–54
- Gardner, R. 2005: 'A Roman cemetery in Clarence Street, Leicester', *Leicestershire Archaeological and Historical Society Transactions* 79, 27–90
- Garnsey, P. 1991: 'Child rearing in ancient Italy', in D.I. Kertzer and R.P. Saller (eds), *The Family in Italy: From Antiquity to the Present*, New Haven, 48–65
- Garnsey, P., and Saller, R. 1987: *The Roman Empire: Economy, Society and Culture*, Berkeley
- Gerrard, J. 2011: 'New radiocarbon dates from the cemetery at Bradley Hill, Somerton', *Proceedings of the Somerset Archaeological and Natural History Society* 154, 189–93
- Goodman, A.H., and Rose, J.C. 1990: 'Assessment of systemic physiological perturbations from dental enamel hypoplasias and associated histological structures', *American Journal of Physical Anthropology* 33 S11, 59–110
- Goodman, A.H., and Rose J.C. 1991: 'Dental enamel hypoplasias as indicators of nutritional status', in M.A. Kelley and C. Larsen (eds), *Advances in Dental Anthropology*, New York, 279–93
- Gowans, E., and Pouncett, J. 2000: *Huntsman's Quarry (Phases 5–7): Excavations and Watching Briefs at Huntsman's Quarry, Naunton, Gloucestershire, 1994–1996*, Gloucester
- Gowland, R., and Garnsey, P. 2010: 'Skeletal evidence for health, nutritional status and malaria in Rome and the empire', in H. Eckardt (ed.), *Roman Diasporas: Archaeological Approaches to Mobility and Diversity in the Roman Empire*, Journal of Roman Archaeology Supplementary Series 78, Portsmouth, RI, 131–56
- Gowland, R., and Redfern, R. 2010: 'Childhood health in the Roman world: perspectives from the centre and margin of the Empire', *Childhood in the Past* 3, 15–42
- Gowland, R., Chamberlain, A., and Redfern, R.C. 2014: 'On the brink of being: re-evaluating infanticide and infant burial in Roman Britain', in Carroll and Graham 2014, 69–88

Grange, J.M. 2001: 'Mycobacterium bovis infection in human beings', *Tuberculosis* 81, 71–77

Halcrow, S.E., and Tayles, N. 2008: 'Stress near the start of life? Localised enamel hypoplasia of the primary canine in late prehistoric mainland Southeast Asia', *Journal of Archaeological Science* 35, 2215–22

Halcrow, S.E., Harris, N.J., Beavan, N., and Buckley, H.R. 2014: 'First bioarchaeological evidence of probable scurvy in Southeast Asia: multifactorial etiologies of vitamin C deficiency in a tropical environment', *International Journal of Palaeopathology* 5, 63–71

Hall, J. 2005: 'The shop-keepers and craft-workers of Roman London', in MacMahon and Price 2005, 125–44

Harinayaran, C.V., Ramalakshmi, T., Prasad, U.V., Sudhakar, D., Srinivasarao, P.V., Sarma, K.V.S., and Kumar, E.G.T. 2007: 'High prevalence of low dietary calcium, high phytate consumption, and vitamin D deficiency in healthy south Indians', *American Journal of Clinical Nutrition* 85, 1062–7

Harlow, M., and Laurence, R. 2002: *Growing Up and Growing Old in Ancient Rome: A Life Course Approach*, London

Harlow, M., and Laurence, R. (eds) 2007: *Age and Ageing in the Roman Empire*, Journal of Roman Archaeology Supplementary Series 65, Portsmouth, RI

Harlow, M., Laurence, R., and Vuolanto, V. 2007: 'Past, present and future in the study of Roman childhood', in S. Crawford and G. Shepherd (eds), *Children, Childhood and Society*, BAR International Series 1696, Oxford, 5–14

Hatton, R.C., and Wall, W. 2006: 'A late Roman cemetery at *Durobrivae*, Chesterton', *Proceedings of the Cambridge Antiquarian Society* 95, 5–24

Heighway, C.M. 1980: 'Roman cemeteries in Gloucester District', *Transactions of the Bristol and Gloucestershire Archaeological Society* 98, 57–72

Hewitt, I. 2012: 'Dewlish Roman villa: post-excavation report', *Proceedings of the Dorset Natural History and Archaeological Society* 133, 65–6

Hillson, S. 1996: *Dental Anthropology*, Cambridge

- Holick, M.F. 2006: 'Resurrection of vitamin D deficiency and rickets', *Journal of Clinical Investigation* 116, 2062–72
- Holley, A.D., Osland, E., Barnes, J., Krishnan, A., and Fraser J.F. 2011: 'Scurvy: historically a plague of the sailor that remains a consideration in the modern intensive care unit', *Internal Medicine Journal* 41, 283–5
- Huck, P. 1995: 'Infant mortality and living standards of English workers during the industrial revolution', *Journal of Economic History* 55, 528–50
- Humphrey, L., Bello, S., and Rousham, E. 2012: 'Sex differences in infant mortality in Spitalfields, London, 1750–1839', *Journal of Biosocial Science* 44, 95–119
- Hurst, H.R. 1985: *Kingsholm: Excavations at Kingsholm Close and Other Sites with a Discussion of the Archaeology of the Area*, Stroud
- Hurst, H.R. 1986: *Gloucester: The Roman and Later Defences*, Stroud
- Jones, A.H.M. 1964: *The Later Roman Empire 284–602: A Social Economic, and Administrative Survey. Volume II*, Baltimore
- Jones, A.H.M. 1974: *The Roman Economy: Studies in Ancient Economic and Administrative History*, Oxford
- Jones, M. 1982: 'Crop production in Roman Britain', in D. Miles (ed.), *The Romano-British Countryside: Studies in Rural Settlement and Economy*, BAR British Series 103, Oxford, 97–108
- Jones, M.E. 1996: *The End of Roman Britain*, London
- Jones, M.J. 2004: 'Cities and urban lives', in Todd 2004, 162–93
- Karl, R. 2005: 'Master and apprentice, knight and squire: education in the 'Celtic' Iron Age', *Oxford Journal of Archaeology* 24, 255–71
- Katzenberg, M.A., Herring, D.A., and Saunders, S.R. 1996: 'Weaning and infant mortality: evaluating the skeletal evidence', *Yearbook of Physical Anthropology* 39, 177–99
- King, A.C. 1984: 'Animal bones and the dietary identity of military and civilian groups in Roman Britain, Germany and Gaul', in T. Blagg and A. King (eds), *Military and Civilian in Roman Britain: Cultural Relationships in a Frontier Province*, BAR British Series 136, Oxford, 187–217

- King, A.C. 1999: 'Diet in the Roman world: a regional inter-site comparison of the mammal bones', *Journal of Roman Archaeology* 12, 168–202
- King, A.C. 2001: 'The Romanization of diet in the Western Empire: comparative archaeological studies', in S. Keay and N. Terrenato (eds), *Italy and the West: Comparative Issues in Romanization*, Oxford, 210–26
- Klaus, H.D. 2012: 'The bioarchaeology of structural violence – a theoretical model and a case study', in D.I. Martin, R.P. Harrod and V.R. Pérez (eds), *The Bioarchaeology of Violence*, Gainesville, 29–62
- Komlos, J. 1998: 'Shrinking in a growing economy? The mystery of physical stature during the industrial revolution', *Journal of Economic History* 58, 779–802
- Kutluk, G., Feyzullah, C., and Muzaffer, B. 2002: 'Comparisons of oral calcium, high dose vitamin D and a combination of these in the treatment of nutritional rickets in children', *Journal of Tropical Pediatrics* 48, 351–3
- Kwon, D.S., Spevak, M.R., Fletcher, K., and Kleinman, P.K. 2002: 'Physiologic subperiosteal new bone formation: prevalence, distribution, and thickness in neonates and infants', *American Journal of Roentgenology* 179, 985–8
- Laes, C. 2004: 'Children and accidents in Roman Antiquity', *Ancient Society* 34, 153–70
- Laes, C. 2007: 'Inscriptions from Rome and the history of childhood', in Harlow and Laurence 2007, 25–36
- Laes, C. 2011: *Children in the Roman Empire: Outsiders Within*, Cambridge
- Lagia, A., Eliopoulos, C., and Manolis, S. 2007: 'Thalassaemia: macroscopic and radiological study of a case', *International Journal of Osteoarchaeology* 17, 269–85
- Lahner, E., Persechino, S., and Annibale, B. 2012: 'Micronutrients (other than iron) and *Helicobacter pylori* infection: a systematic review', *Helicobacter* 17, 1–15
- Laurence, R., Esmonde Cleary, S., and Sears, G. 2011: *The City in the Roman West c.250 BC – c.AD 250*, Cambridge
- Lawson, J.P., Ablow, R.C., and Pearson, H.A. 1981a: 'The ribs in thalassemia. I. The relationship to therapy', *Radiology* 140, 663–72

- Lawson, J.P., Ablow, R.C., and Pearson, H.A. 1981b: 'The ribs in thalassaemia. II. The pathogenesis of the changes', *Radiology* 140, 673–9
- Leech, R. 1982: *Excavations at Catsgore 1970–1973: A Romano-British Village*, Western Archaeological Trust Excavation Monograph 2, Taunton
- Leech, R., Besly, E.M., and Everton, R.F. 1981: 'The excavation of a Romano-British farmstead and cemetery on Bradley Hill, Somerton, Somerset', *Britannia* 12, 177–252
- Lewis, M.E. 2004: 'Endocranial lesions in non-adult skeletons: understanding their aetiology', *International Journal of Osteoarchaeology* 14, 82–97
- Lewis, M.E. 2007: *The Bioarchaeology of Children: Perspectives from Biological and Forensic Anthropology*, Cambridge
- Lewis, M.E. 2010: 'Life and death in a civitas capital: metabolic disease and trauma in the children from late Roman Dorchester, Dorset', *American Journal of Physical Anthropology* 142, 405–16
- Lewis, M.E. 2011: 'Tuberculosis in the non-adults from Romano-British Poundbury Camp, Dorset, England', *International Journal of Palaeopathology* 1, 12–23
- Lewis, M.E. 2012: 'Thalassaemia: its diagnosis and interpretation in past skeletal populations', *International Journal of Osteoarchaeology* 22, 685–93
- Lewis, M.E., and Gowland, R. 2007: 'Brief and precarious lives: infant mortality in contrasting sites from medieval and post-medieval England (AD 850–1859)', *American Journal of Physical Anthropology* 134, 117–29
- Lewis, M.E., and Roberts, C. 1997: 'Growing pains: the interpretation of stress indicators', *International Journal of Osteoarchaeology* 7, 581–6
- Lovell, N.C. 1997: 'Trauma analysis in paleopathology', *Yearbook of Physical Anthropology* 40, 139–70
- Mander, J. 2013: *Portraits of Children on Roman Funerary Monuments*, Cambridge
- Marais, B.J. 2011: 'Childhood tuberculosis: epidemiology and natural history of disease', *Indian Journal of Pediatrics* 78, 321–7
- Matthews, C.L. 1981: 'A Romano-British inhumation cemetery at Dunstable', *Bedfordshire Archaeological Journal* 15

- Mattingly, D. 2006: *An Imperial Possession: Britain in the Roman Empire*, London
- Mays, S. 2008: 'Metabolic bone disease', in R. Pinhasi and S. Mays (eds), *Advances in Human Palaeopathology*, Chichester, 215–52
- McCarthy, M. 2013: *The Romano-British Peasant: Towards a Study of People, Landscapes and Work during the Roman Occupation of Britain*, Oxford
- McConnell, D., Grassam, A., and Mustchin, A. 2012: *Land Adjacent to Great Casterton Primary School, Pickworth Road, Great Casterton, Rutland, Suffolk*
- McDade, T.W., and Worthman, C. M. 1998: 'The weanling's dilemma reconsidered: a biocultural analysis of breastfeeding ecology', *Journal of Developmental and Behavioural Pediatrics* 19, 286–99
- McIlvaine, B.K. 2013: 'Implications of reappraising the iron-deficiency anemia hypothesis', *International Journal of Osteoarchaeology* 25, 997–1000
- MacMahon, A. 2005: 'The shops and workshops of Roman Britain', in MacMahon and Price 2005, 48–69
- MacMahon, A., and Price, J. (eds) 2005: *Roman Working Lives and Urban Living*, Oxford
- McWilliam, J. 2001: 'Children among the dead – the influence of urban life on the commemoration of children on tombstone inscriptions', in S. Dixon (ed.), *Childhood, Class and Kin in the Roman World*, London, 74–98
- Mensforth, R.P., Lovejoy, C.O., Lallo, J.W., and Armelagos, G.J. 1978: 'The role of constitutional factors, diet, and infectious disease in the etiology of porotic hyperostosis and periosteal reactions in prehistoric infants and children', *Medical Anthropology* 2, 1–59
- Millett, M. 1999: 'Coloniae and Romano-British studies', in H. Hurst (ed.), *The Coloniae of Roman Britain: New Studies and a Review*, Journal of Roman Archaeology Supplementary Series 36, Portsmouth, RI, 191–6
- Millett, M. 2001: 'Approaches to urban societies', in S. James and M. Millett (eds), *Britons and Romans: Advancing an Archaeological Agenda*, Council for British Archaeology Research Report 125, York, 60–6
- Millett, M. 2005: *Roman Britain*, London

- Millett, M., and Gowland, R. 2015: 'Infant and child burial rites in Roman Britain: a study from East Yorkshire', *Britannia* 46, 171–89
- Mirković, M. 1997: 'The later Roman colonate and freedom', *Transactions American Philosophical Society* 87, 1–144
- Molleson, T. 1992: 'The anthropological evidence for change through Romanisation of the Poundbury population', *Anthropologischer Anzeiger* 50, 179–89
- Moore, A. 2009: 'Hearth and home: the burial of infants within Romano-British domestic contexts', *Childhood in the Past* 2, 33–54
- Moorrees, C.F.A., Fanning, E.A., and Hunt, E.E. 1963a: 'Age variation of formation stages for ten permanent teeth', *Journal of Dental Research* 42, 1490–502
- Moorrees, C.F.A., Fanning, E.A., and Hunt, E.E. 1963b: 'Formation and resorption of three deciduous teeth in children', *American Journal of Physical Anthropology* 21, 205–13
- Mouritsen, H. 2011: 'The families of Roman slaves and freedmen', in B. Rawson (ed.), *A Companion to Families in the Greek and Roman Worlds*, Chichester, 129–44
- Mullin, D. 2006: *The Archaeological Landscape of Frampton-on-Severn, Gloucestershire: Updated Project Design*, Gloucester
- Müldner, G. 2013: 'Stable isotopes and diet: their contribution to Romano-British research', *Antiquity* 87, 137–49
- Nehlich, O., Fuller, B.T., Jay, M., Mora, A., Nicholson, R.A., Smith, C.I., and Richards, M.P. 2011: 'Application of sulphur isotope ratios to examine weaning patterns and freshwater fish consumption in Roman Oxfordshire, UK', *Geochimica et Cosmochimica Acta* 75, 4963–77
- Nelson, L.J., and Wells, C.D. 2004: 'Global epidemiology of childhood tuberculosis', *International Journal of Tuberculosis and Lung Disease* 8, 636–47
- Nicholson, K. 2006: 'A late Roman cemetery at Watersmeet, Mill Common, Huntingdon', *Proceedings of the Cambridge Antiquarian Society* 95, 57–90
- Nielsen, A.V.F., Tetens, I., and Meyer, A.S. 2013: 'Potential of phytase-mediated iron release from cereal-based foods: a quantitative view', *Nutrients* 5, 3074–98

- Norman, N.J. 2002: 'Death and burial of Roman children: the case of the Yasmina cemetery at Carthage – Part I, setting the stage', *Mortality* 7, 302–23
- Ogden, J.A. 2000: *Skeletal Injury in the Child* (3rd edn), New York
- Ortner, D.J. 2003: *Identification of Pathological Conditions in Human Skeletal Remains* (2nd edn), San Diego
- Ortner, D.J., and Ericksen, M.F. 1997: 'Bone changes in the human skull probably resulting from scurvy in infancy and childhood', *International Journal of Osteoarchaeology* 7, 212–20
- Ortner, D.J., and Mays, S. 1998: 'Dry-bone manifestations of rickets in infancy and early Childhood', *International Journal of Osteoarchaeology* 8, 45–55
- Ortner, D., Butler, W., Cafarella, J., and Milligan, L. 2001: 'Evidence of probable scurvy in subadults from archaeological sites in North America', *American Journal of Physical Anthropology* 114, 343–51
- Ortner, D., Kimmerle, E., and Diez, M. 1999: 'Probable evidence of scurvy in subadults from archaeological sites in Peru', *American Journal of Physical Anthropology* 108, 321–31
- Ottaway, P. 2009: *Roman York*, Stroud
- Ottaway, P.J., Qualmann, K.E., Rees, H., and Scobie, G.D. 2012: *The Roman Cemeteries and Suburbs of Winchester: Excavations 1971–86*, Winchester
- Oxenham, M.F., and Cavill, I. 2010: 'Porotic hyperostosis and cribra orbitalia: the erythropoietic response to iron-deficiency anaemia', *Anthropological Science* 118, 199–200
- Pearce, S.M. 1982: 'Estates and church sites in Dorset and Gloucestershire: the emergence of a Christian society', in S.M. Pearce (ed.), *The Early Church in Western Britain and Ireland: Studies Presented to C. A. Raleigh Radford*, BAR British Series 102, Oxford, 117–37
- Pearce, J. 1999a: *Case Studies in a Contextual Archaeology of Burial Practice in Roman Britain*, unpub. PhD Thesis, University of Durham
- Pearce, J. 1999b: 'The dispersed dead: preliminary observations on burial and settlement space in rural Roman Britain', in P. Baker, C. Forcey, S. Jundi and R. Witcher (eds),

- TRAC 98: *Proceedings of the Eighth Annual Theoretical Roman Archaeology Conference, Leicester 1998*, Oxford, 151–62
- Pearce, J. 2001: 'Infants, cemeteries and communities in the Roman provinces', in D. Davies, A. Gardner and K. Lockyear (eds), *TRAC 2000: Proceedings of the Tenth Annual Theoretical Roman Archaeology Conference, London 2000*, Oxford, 125–42
- Pearce, J. 2015: 'Urban exits: commercial archaeology and the study of death rituals and the dead in the towns of Roman Britain', in M. Fulford and N. Holbrook (eds), *The Towns of Roman Britain: The Contribution of Commercial Archaeology Since 1990*, Britannia Monograph Series 27, Cambridge, 138–66
- Percival, J. 1976: *The Roman Villa*, London
- Pettifor, J.M. 2014: 'Calcium and vitamin D metabolism in children in developing countries', *Annals of Nutrition and Metabolism* 64, 15–22
- Pfeiffer, S. 1984: 'Paleopathology in an Iroquoian ossuary, with special reference to tuberculosis', *American Journal of Physical Anthropology* 65, 181–9
- Pfeiffer, S. 1991: 'Rib lesions and New World tuberculosis', *International Journal of Osteoarchaeology* 1, 191–8
- Philpott, R. 1991: *Burial Practice in Roman Britain: A Survey of Grave Treatment and Furnishing AD 43–410*, BAR British Series 219, Oxford
- Pitts, M., and Griffin, R. 2012: 'Exploring health and social well-being in late Roman Britain: an intercemetary approach', *American Journal of Archaeology* 116, 253–76
- Pliny the Elder: *Natural History Volume II: Books 3–7*, Loeb Classical Library 352, translated by H. Rackham (1942), Cambridge, Mass./London
- Popovich, D., McAlhany, A., Adewumi, A.O., and Barnes, M.M. 2009: 'Scurvy: forgotten but definitely not gone', *Journal of Pediatric Health Care* 23, 405–15
- Powell, L.A., Redfern, R.C., Millard, A.R., and Gröcke, D.R. 2014: 'Infant feeding practices in Roman London: evidence from isotopic analyses', in Carroll and Graham 2014, 89–110
- Pre-Construct Archaeology. 2006: *Bantycok Gypsum Mine, Nottinghamshire*, Lewisham
- Price, E. 2000a: *Frocester: A Romano-British Settlement, its Antecedents and Successors. Volume 1: The Sites*, Stonehouse

- Price, E. 2000b: *Frocester: A Romano-British Settlement, its Antecedents and Successors. Volume 2: The Finds*, Stonehouse
- Putnam, B. 2007: *Roman Dorset*, Stroud
- Rahtz, P., Hirst, S., and Wright, S.M. 2000: *Cannington Cemetery: Excavations 1962–3 of Prehistoric, Roman, post-Roman, and later features at Cannington Park Quarry, near Bridgwater, Somerset*, Britannia Monograph Series 17, Cambridge
- Rana, R.S., Wu, J.S., and Eisenberg, R.L. 2009: 'Periosteal reaction', *American Journal of Roentgenology* 193, 259–72
- Rawson, B. 2003a: *Children and Childhood in Roman Italy*, Oxford
- Rawson, B. 2003b: 'The Roman family in recent research: state of the question', *Biblical Interpretation* 11, 119–38
- Redfern, R. 2007: 'The influence of culture upon childhood: an osteological study of Iron Age and Romano-British Dorset, England', in Harlow and Laurence 2007, 171–90
- Redfern, R., and Roberts, C.A. 2005: 'Health in Romano-British urban communities: reflections from the cemeteries', in D.N. Smith, M.B. Brickley and W. Smith (eds), *Fertile Ground: Papers in Honour of Susan Limbrey*, Oxford, 115–29
- Redfern, R.C., Millard, A.R., and Hamlin, C. 2012: 'A regional investigation of subadult dietary patterns and health in late Iron Age and Roman Dorset', *Journal of Archaeological Science* 39, 1249–59
- Redfern, R.C., DeWitte, S.N., Pearce, J., Hamlin, C., and Dinwiddy, K.E. 2015: 'Urban-rural differences in Roman Dorset, England: a bioarchaeological perspective on Roman settlements', *American Journal of Physical Anthropology* 157, 107–20
- Resnick, D., and Kransdorf, M.J. 2005: *Bone and Joint Imaging* (3rd edn), Philadelphia
- Revell, L. 2005: 'The Roman life course: a view from the inscriptions', *European Journal of Archaeology* 8, 43–63
- Roberts, C.A. 2007: 'A bioarchaeological study of maxillary sinusitis', *American Journal of Physical Anthropology* 133, 792–807
- Roberts, C.A., and Buikstra, J.E. 2003: *The Bioarchaeology of Tuberculosis: A Global View on a Re-emerging Disease*, Gainesville

- Roberts, C., and Cox, M. 2003: *Health and Disease in Britain: From Prehistory to the Present Day*, Stroud
- Roberts, C., and Manchester, K. 2010: *The Archaeology of Disease* (3rd edn), Stroud
- Robit, B., Aranda, N., Giralt, M., Romen, M., Balaguer, A., and Arijia, V. 2013: 'Effect of different doses of iron supplementation during pregnancy on maternal and infant health', *Annals of Hematology* 92, 221–9
- Rohnbogner, A. 2015: *Dying Young. A Palaeopathological Study of Child Health in Roman Britain*, unpub. PhD Thesis, University of Reading
- Rohnbogner, A. 2016: 'Differential diagnosis of a probable case of non-adult thalassaemia from 4th century AD Romano-British Colchester, UK', *International Journal of Palaeopathology*, doi: 10.1016/j.ijpp.2016.08.002
- Rohnbogner, A., and Lewis, M.E. 2016: 'Poundbury Camp in context – a new perspective on the lives of children from urban and rural Roman England', *American Journal of Physical Anthropology*, doi: 10.1002/ajpa.23106
- Rohnbogner, A., and Lewis, M.E. forthcoming: 'Dental caries as a measure of diet, health and difference in non-adults from urban and rural Roman Britain', *Dental Anthropology*
- Roksandic, M. 2003: 'New standardised visual forms for recording the presence of human skeletal elements in archaeological and forensic contexts', *Internet Archaeology* 01/2003
- Sallares, R. 2002: *Malaria and Rome: A History of Malaria in Ancient Italy*, Oxford
- Scheidel, W., and von Reden, S. (eds) 2002: *The Ancient Economy*, New York
- Scheuer, L., and Black, S. 2000: *Developmental Juvenile Osteology*, San Diego
- Scheuer, J.L., Musgrave, J.H., and Evans, P. 1980: 'The estimation of fetal and perinatal age from limb bone length by linear and logarithmic regression', *Annals of Human Biology* 7, 257–65
- Schultz, M. 2003: 'Light microscopic analysis in skeletal palaeopathology', in D.J. Ortner, *Identification of Pathological Conditions in Human Skeletal Remains* (2nd edn), London, 73–107
- Scott, E. 1991: 'Animal and infant burials in Romano-British villas: a revitalization movement', in P. Garwood (ed.), *Sacred and Profane: Proceedings of a Conference on*

- Archaeology, Ritual and Religion, Oxford*, Oxford University Committee for Archaeology Monograph, Oxford, 115–21
- Scott, S., and Duncan, C.J. 1999: 'Malnutrition, pregnancy and infant mortality: a biometric model', *Journal of Interdisciplinary History* 30, 37–60
- Shin, J.S., Choi, M.Y., Longtine, M.S., and Nelson, D.M. 2010: 'Vitamin D effects on pregnancy and the placenta', *Placenta* 31, 1027–34
- Skinner, M.F., and Hung, J.T.W. 1989: 'Social and biological correlates of localized enamel hypoplasia of the human deciduous canine tooth', *American Journal of Physical Anthropology* 79, 159–75
- Smith, R.J.C., Healy, F., Allen, M.J., Morris, E.L., Barnes, I., and Woodward, P.J. 1997: *Excavations along the Route of the Dorchester By-pass, Dorset, 1986–8*, Wessex Archaeology Report 11, Salisbury
- Soren, D. 2003: 'Can archaeologists excavate evidence of malaria?', *World Archaeology* 35, 193–209
- Soranus: *Soranus' Gynecology* (trans. O. Temkin, 1991), Baltimore
- Stark, R.J. 2014: 'A proposed framework for the study of paleopathological cases of subadult scurvy', *International Journal of Palaeopathology* 5, 18–26
- Stead, I.M., and Rigby, V. 1986: *Baldock: The Excavation of a Roman and Pre-Roman Settlement, 1968–72*, London
- Stead, W.W. 2000: 'What's in a name: confusion of *Mycobacterium tuberculosis* and *Mycobacterium bovis* in ancient DNA analysis', *Palaeopathology Association Newsletter* 110, 13–16
- Stuart-Macadam, P. 1985: 'Porotic hyperostosis: representative of a childhood condition', *American Journal of Physical Anthropology* 66, 391–8
- Stuart-Macadam, P. 1991: 'Anaemia in Roman Britain: Poundbury Camp', in H. Bush and M. Zvelebil (eds), *Health in Past Societies: Biocultural Interpretations of Human Skeletal Remains in Archaeological Contexts*, Oxford, 101–13

- Thacher, T.D., Fischer, P.R., Strand, M.A., and Pettifor, J.M. 2006: 'Nutritional rickets around the world: causes and future directions', *Paediatrics and International Child Health* 26, 1–16
- Timberlake, S., Dodwell, N., and Armour, N. 2007: *The Roman Cemetery, The Babraham Institute, Cambridgeshire: An Archaeological Excavation*, Cambridge Archaeological Unit Report 754, Cambridge
- Todd, M. 1975: 'Margidunum and Ancaster', in W. Rodwell and T. Rowley (eds), *The 'Small Towns' of Roman Britain*, British Archaeological Report 15, Oxford, 211–24
- Todd, M. (ed.) 2004: *A Companion to Roman Britain*, Oxford
- Tomlin, R.S.O. 2003: "'The girl in question': a new text from Roman London", *Britannia* 34, 41–51
- Tyler, P.A., Madani, G., Chaudhuri, R., Wilson, L.F., and Dick, E.A. 2006: 'The radiological appearances of thalassaemia', *Clinical Radiology* 61, 40–2
- USDA 2013: *USDA National Nutrient Database for Standard Reference Release 26 – Nutrients: Vitamin C, Total Ascorbic Acid (mg)*, Available from: <http://ods.od.nih.gov> [accessed 07/05/2015]
- van der Veen, M. 2007: 'Food as an instrument of social change: feasting in Iron age and early Roman southern Britain', in K.C. Twiss (ed.), *The Archaeology of Food and Identity*, Southern Illinois University Carbondale Occasional Paper 34, Carbondale, 112–29
- van der Veen, M. 2008: 'Food as embodied material culture: diversity and change in plant food consumption in Roman Britain', *Journal of Roman Archaeology* 21, 83–109
- van der Veen, M., Livarda, A., and Hill, A. 2007: 'The archaeobotany of Roman Britain: current state and identification of research priorities', *Britannia* 38, 181–210
- van der Veen, M., Livarda, A., and Hill, A. 2008: 'New plant foods in Roman Britain: dispersal and social access', *Environmental Archaeology* 13, 11–36
- van Gerven, D.P., Sanford, M.K., and Hummert, J.R. 1981: 'Mortality and culture change in Nubia's Batn el Hajar', *Journal of Human Evolution* 10, 395–408

- van Rijn, R.R., Bilo, R.A.C., and Robben, S.G.F. 2009: 'Birth-related, mid-posterior rib fractures in neonates: a report of three cases (and a possible fourth case) and a review of the literature', *Pediatric Radiology* 39, 30–4
- van Sleuwen, B.E., Engelberts, A.C., Boere-Boonekamp, M.M., Kuis, W., Schulpen, T.W. J., and L'Hoir, M.P. 2007: 'Swaddling: a systematic review', *Pediatrics* 120, 1097–106
- Verlinden, P., and Lewis, M.E. 2015: 'Childhood trauma: methods for the identification of physeal fractures in nonadult skeletal remains', *American Journal of Physical Anthropology* 157, 411–20
- Viner, L., and Leech, R. 1982: 'Bath Gate Cemetery, 1969–1976', in A. McWhirr, L. Viner and C. Wells, *Cirencester Excavations II: Romano-British Cemeteries at Cirencester*, Cirencester, 69–133
- Vögele, J.P. 1994: 'Urban infant mortality in Imperial Germany', *Social History of Medicine* 7, 401–26
- Walker, P. 1997: 'Skeletal evidence for child abuse: a physical anthropological perspective', *Journal of Forensic Sciences* 42, 196–207
- Walker, P.L., Bathurst, R.R., Richman, R., Gjerdrum, T., and Andrushko, V.A. 2009: 'The causes of porotic hyperostosis and cribra orbitalia: a reappraisal of the iron-deficiency-anemia hypothesis', *American Journal of Physical Anthropology* 139, 109–25
- Walls, T., and Shingadia, D. 2004: 'Global epidemiology of pediatric tuberculosis', *Journal of Infection* 48, 13–22
- Wapler, U., Cruzeby, E., Schultz, M. 2004: 'Is cribra orbitalia synonymous with anemia? Analysis and interpretation of cranial pathology in Sudan', *American Journal of Physical Anthropology* 123, 333–9
- Webster, J. 2005: 'Archaeologies of slavery and servitude: bringing 'New World' perspectives to Roman Britain', *Journal of Roman Archaeology* 18, 161–79
- Wenham, L.P. 1968: *The Romano-British Cemetery at Trentholme Drive, York*, Ministry of Public Building and Works Archaeological Reports 5, London
- Weston, D.A., 2008: 'Investigating the specificity of periosteal reactions in pathology museum specimens', *American Journal of Physical Anthropology* 137, 48–59

- Whittaker, C.R., and Garnsey, P. 1997: 'Rural life in the later Roman Empire', in A. Cameron and P. Garnsey (eds), *The Cambridge Ancient History 13. The Late Empire, AD 337–425*, Cambridge, 277–311
- Wileman, J. 2005: *Hide and Seek: The Archaeology of Childhood*, Stroud
- Wood, J.W., Milner, G.R., Harpending, H.C., and Weiss, K.M. 1992: 'The osteological paradox – problems of inferring prehistoric health from skeletal samples', *Current Anthropology* 33, 343–70
- WORD 2016: *Roman Data*, Museum of London [Accessed 10/09/2016]
- Yurdakok, K., Yavuz, T., and Taylor, C.E. 1990: 'Swaddling and acute respiratory infections', *American Journal of Public Health* 80, 873–5

¹ Roberts and Cox 2003, ch. 3.

² Pitts and Griffin 2012; Redfern *et al.* 2015.

³ Fuller *et al.* 2006; Nehlich *et al.* 2011; Redfern *et al.* 2012; Powell *et al.* 2014.

⁴ Mensforth *et al.* 1978; Lewis 2007, 19.

⁵ Gowland and Redfern 2010.

⁶ e.g. Rawson 2003a; 2003b; McWilliam 2001; Harlow and Laurence 2002; Revell 2005; Harlow *et al.* 2007; Laes 2004; 2007; Crummy 2010; Mander 2013.

⁷ Bradley 1984, 77–9; Garnsey 1991; Garnsey and Saller 1987, 108–20; Pearce 2001; Laes 2007, 25; Mouritsen 2011, 129.

⁸ Burn 1970; McWilliam 2001; Tomlin 2003; Revell 2005; Laes 2007.

⁹ Soranus, *Gynaecology* II.

¹⁰ *ibid.* II.21.

¹¹ Gowland and Garnsey 2010.

¹² Ayi *et al.* 2004.

¹³ Harinarayan *et al.* 2007; Walker *et al.* 2009; Oxenham and Cavill 2010.

¹⁴ Sallares 2002; Soren 2003.

¹⁵ Gowland and Garnsey 2010.

¹⁶ Redfern 2007; Redfern *et al.* 2012.

¹⁷ Lewis 2010; 2011; 2012.

¹⁸ Lewis 2010.

¹⁹ Lewis 2012.

²⁰ Cool 2006.

²¹ Lewis 2010; 2012.

²² Soranus, *Gynaecology* II.21; Lewis 2010.

²³ Yurdakok *et al.* 1990; Kutluk *et al.* 2002; van Sleuwen *et al.* 2007.

²⁴ Soranus, *Gynaecology* II.21.

²⁵ Lewis 2011.

²⁶ Stead 2000; Roberts and Buikstra 2003; Roberts and Manchester 2010, 187.

²⁷ See Rohnbogner and Lewis 2016 where these results are contextualised using both urban and rural contemporary non-adult samples.

²⁸ Webster 2005; Mattingley 2006; McCarthy 2013.

²⁹ Pitts and Griffin 2012.

³⁰ Redfern *et al.* 2015.

³¹ WORD 2016.

³² Rohnbogner and Lewis 2016.

-
- ³³ Millett 1999; 2001; 2005; Mattingly 2006; Laurence *et al.* 2011; Pitts and Griffin 2012; McCarthy 2013, Breeze 2014; Pearce 2015.
- ³⁴ Moorrees *et al.* 1963a; 1963b.
- ³⁵ Scheuer *et al.* 1980.
- ³⁶ Scheuer and Black 2000.
- ³⁷ Walker *et al.* 2009.
- ³⁸ Mensforth *et al.* 1978; Stuart-Macadam 1985.
- ³⁹ Walker *et al.* 2009; Oxenham and Cavill 2010; McIlvaine 2013
- ⁴⁰ Lewis 2012.
- ⁴¹ Hillson 1996, 165.
- ⁴² Goodman and Rose 1990; 1991.
- ⁴³ Goodman and Rose 1990; 1991; Roberts and Manchester 2010, 75.
- ⁴⁴ Skinner and Hung 1989; Halcrow and Tayles 2008.
- ⁴⁵ Lewis and Roberts 1997; Lewis 2007, 141.
- ⁴⁶ Schultz 2003, 93; Lewis 2004.
- ⁴⁷ Walker 1997; Lewis 2007, 134–5.
- ⁴⁸ Weston 2008.
- ⁴⁹ Roberts and Manchester 2010, 187.
- ⁵⁰ Marais 2011.
- ⁵¹ Stead 2000; Grange 2011.
- ⁵² Holick 2006.
- ⁵³ Thacher *et al.* 2006; Shin *et al.* 2010; Pettifor 2014.
- ⁵⁴ Mays 2008, 223.
- ⁵⁵ Ortner and Ericksen 1997; Brickley and Ives 2006.
- ⁵⁶ Armelagos *et al.* 2014; Buckley *et al.* 2014; Crandall 2014; Crandall and Klaus 2014; Halcrow *et al.* 2014; Stark 2014.
- ⁵⁷ Ortner 2003, ch. 8.
- ⁵⁸ Verlinden and Lewis 2015.
- ⁵⁹ Skinner and Hung 1989; Goodman and Rose 1990.
- ⁶⁰ After Lewis and Roberts 1997; Ortner 2003, 206–14; Lewis 2004; 2007, 134–43; Weston 2008.
- ⁶¹ Pfeiffer 1984; 1991; Roberts and Buikstra 2003; Resnick and Kransdorf 2005, 758–63.
- ⁶² Stuart-Macadam 1991.
- ⁶³ Lawson *et al.* 1981a; 1981b.
- ⁶⁴ Ortner 2003, 364–6; Tyler *et al.* 2006; Lagia *et al.* 2007; Lewis 2011.
- ⁶⁵ Ortner and Mays 1998; Ortner 2003, 393–404; Brickley and Ives 2008.
- ⁶⁶ Brickley and Ives 2006.
- ⁶⁷ Ortner and Ericksen 1997; Ortner *et al.* 1999; 2001; Crandall and Klaus 2014; Stark 2014.
- ⁶⁸ See Rohnbogner 2016.
- ⁶⁹ Kwon *et al.* 2002; Lewis 2004; 2007, 135–6; Rana *et al.* 2009; also see Brickley and Ives 2006.
- ⁷⁰ Clough and Boyle 2010, 386–9; see Rohnbogner and Lewis, forthcoming.
- ⁷¹ See Rohnbogner 2015 for case studies.
- ⁷² Clough and Boyle 2010, 390.
- ⁷³ Also see Rohnbogner 2016 for a case of a radial fracture in a 1.1–2.5 year old with suspected thalassaemia and active vitamin D deficiency.
- ⁷⁴ Ogden 2000, 431, 466, 596.
- ⁷⁵ *ibid.*, 880, 1035.
- ⁷⁶ Barsness *et al.* 2003.
- ⁷⁷ Bulloch *et al.* 2000; van Rijn *et al.* 2009.
- ⁷⁸ Lovell 1997.
- ⁷⁹ Lewis 2010.
- ⁸⁰ See Rohnbogner and Lewis 2016.
- ⁸¹ Wood *et al.* 1992
- ⁸² Frenzen and Hogan 1982.
- ⁸³ Scott and Duncan 1999.
- ⁸⁴ Vögele 1994; Lewis and Gowland 2007; Humphrey *et al.* 2012.
- ⁸⁵ Rawson 2003a, 121; Roberts and Cox 2003, 123–30.
- ⁸⁶ Gowland *et al.* 2014; Millett and Gowland 2015.
- ⁸⁷ For example the third–fifth century Yasminea necropolis of Carthage, Norman 2002.
- ⁸⁸ Pliny the Elder, *Natural History* VII.16.
- ⁸⁹ Philpott 1991, 101.

-
- ⁹⁰ Philpott 1991, 101; Scott 1991, 120; Pearce 1999a; 1999b; Esmonde Cleary 2000; Wileman 2005, 80–1, 99; Moore 2009; Gowland *et al.* 2014.
- ⁹¹ Thacher *et al.* 2006.
- ⁹² Brickley and Ives 2008; Pettifor 2014.
- ⁹³ Foote and Marriott 2003.
- ⁹⁴ McDade and Worthman 1998; Fewtrell *et al.* 2007.
- ⁹⁵ Katzenberg *et al.* 1996; Robit *et al.* 2013.
- ⁹⁶ Stuart-Macadam 1985; Oxenham and Cavill 2010.
- ⁹⁷ Lewis 2010.
- ⁹⁸ Roberts and Buikstra 2003; Roberts 2007.
- ⁹⁹ Roberts and Manchester 2010, 184.
- ¹⁰⁰ Nelson and Wells 2004; Walls and Shingadia 2004.
- ¹⁰¹ Redfern and Roberts 2005, 122; Gowland and Redfern 2010; Redfern *et al.* 2015.
- ¹⁰² Roberts and Cox 2003, 123.
- ¹⁰³ Karl 2005; Webster 2005.
- ¹⁰⁴ Bradley 1991, 108–10; Rawson 2003a, 121; Laes 2011, 195.
- ¹⁰⁵ Whittaker and Garnsey 1997; Esmonde Cleary 2004; McCarthy 2013, 7, 90.
- ¹⁰⁶ Previously reported by Brothwell *et al.* 2000, 203.
- ¹⁰⁷ Harlow and Laurence, 2002.
- ¹⁰⁸ Rahtz *et al.* 2000, 423; McCarthy 2013, 101–2.
- ¹⁰⁹ Jones 2004; Hall 2005, 137.
- ¹¹⁰ MacMahon 2005; Gowland and Garnsey 2010, 149; Lewis 2010.
- ¹¹¹ Halcrow *et al.* 2014.
- ¹¹² Popovich *et al.* 2009; Holley *et al.* 2011; Lahner *et al.* 2012; Halcrow *et al.* 2014.
- ¹¹³ Crandall and Klaus 2014; Stark 2014.
- ¹¹⁴ Wapler *et al.* 2004; Djuric *et al.* 2008; Walker *et al.* 2009; Oxenham and Cavill 2010.
- ¹¹⁵ King 1984; 1999; 2001; Molleson 1992; van der Veen 2007; 2008; van der Veen *et al.* 2007; 2008; Cummings 2009; Müldner 2013.
- ¹¹⁶ Facchini *et al.* 2004; Nielsen *et al.* 2013.
- ¹¹⁷ USDA 2013.
- ¹¹⁸ Klaus 2012; Crandall 2014; Armelagos *et al.* 2014.
- ¹¹⁹ Huck 1995; Komlos 1998; Lewis and Gowland 2007.
- ¹²⁰ Crandall 2014.
- ¹²¹ van der Veen *et al.* 2008; McCarthy 2013, 58–9; Breeze 2014.
- ¹²² Scheidel and von Reden 2002.
- ¹²³ Jones 1982; Pearce 1982; Faith 1997, 17–18; Whittaker and Garnsey 1997.
- ¹²⁴ Jones 1964, ch. 20; 1974, 293; Percival 1976, 120; Jones 1996, 208–13; Mirković 1997.
- ¹²⁵ Jones 1996, 208–15; Esmonde Cleary 2004.
- ¹²⁶ Whittaker and Garnsey 1997, 309–11.
- ¹²⁷ Jones 1982, 2004; Esmonde Cleary 2004; Fulford 2004, 316.
- ¹²⁸ van Gerven *et al.* 1981; Jones 1996, 215; Whittaker and Garnsey 1997, 285.
- ¹²⁹ Jones 1982; Faith 1997, 16–18; Whittaker and Garnsey 1997, 301; Esmonde Cleary 2004.
- ¹³⁰ Farmer 2008.

Table 1. Individual site assemblage details.

Site	Date (A.D.)	Type	Non-adult n	Site Reference
Winchester (North, West, East)*	1-4 th century	Major Urban	166	Ottaway <i>et al.</i> 2012
Kingsholm, Gloucester*	2-4 th century	Major Urban	17	Hurst 1985; 1986
Gambier-Parry Lodge, Gloucester*	2-4 th century	Major Urban	12	Heighway 1980; Mullin 2006
Trentholme Drive, York*	3-4 th century	Major Urban	24	Wenham 1968; Ottaway 2009
Clarence Street, Leicester	3-4 th century	Major Urban	13	Gardner 2005
Bath Gate, Cirencester*	4 th century	Major Urban	64	Viner and Leech 1982
Lankhills, Winchester	4 th century	Major Urban	67	Booth <i>et al.</i> 2010
London	4 th century	Major Urban	50	WORD 201
Butt Road, Colchester	4-5 th century	Major Urban	109	Crummy and Crossan 1993
Major urban total N			522	
Springhead, Kent	1-4 th century	Minor Urban	82	Boyle and Early 1999; Barnett <i>et al.</i> 2011
Baldock, Hertfordshire*	2-4 th century	Minor Urban	83	Stead and Rigby 1986; Burleigh and Fitzpatrick-Matthews 2010
Queenford Farm/Queensford Mill, Oxfordshire*	3-4 th century	Minor Urban	60	Durham and Rowley 1972; Chambers 1987
Ancaster, Lincolnshire*	3-4 th century	Minor Urban	81	Todd 1975
Great Casterton, Rutland*	3-4 th century	Minor Urban	38	McConnell <i>et al.</i> 2012
Ashton, Northamptonshire*	4 th century	Minor Urban	60	Dix 1983
Dunstable, Bedfordshire*	3-5 th century	Minor Urban	27	Matthews 1981
Chesterton, Cambridgeshire	3-5 th century	Minor Urban	9	Hatton and Wall 2006
Minor urban total N			440	
Owslebury, Hampshire*	1-4 th century	Rural	16	Collis 1968; 1977
Huntsman's Quarry, Gloucestershire	2-3 rd century	Rural	12	Gowans and Pouncett 2000
Babraham Institute, Cambridgeshire	2-4 th century	Rural	12	Timberlake <i>et al.</i> 2007
Bantycok Mine, Nottinghamshire	2-4 th century	Rural	7	Pre-Construct Archaeology 2006
Catsgore, Somerset*	2-5 th century	Rural	19	Leech 1982
Cannington, Somerset	3-4 th century	Rural	148	Rahtz <i>et al.</i> 2000
Dorchester by-pass, Dorset	4 th century	Rural	9	Smith <i>et al.</i> 1997
Dewlish, Dorset	4 th century	Rural	8	Putnam 2007; Hewitt 2012
Bradley Hill, Somerset*#	4-5 th century	Rural	29	Leech <i>et al.</i> 1981; Gerrard 2011
Watersmeet, Cambridgeshire	4-5 th century	Rural	14	Nicholson 2006
Frocester, Gloucestershire	3-5 th century	Rural	43	Price 2000a; 2000b
Rural total N			317	
Total sample N			1,279	

* site samples recorded by the author to attain primary data #non-adult burials were recovered from within the settlement boundaries associated with a fourth-century building (Leech *et al.* 1981), rather than from the fifth-century cemetery (Gerrard 2011).

Table 2. Ages-at-death of non-adults from major urban, minor urban and rural settlements of Roman England.

Age (yrs)	Major urban		Minor urban		Rural		Total	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Perinate	88	16.9	84	19.1	95	29.9	267	20.9
0.0-1.0	83	15.9	145	32.9	77	24.3	305	23.8
1.1-2.5	55	10.5	57	12.9	34	10.7	146	11.4
2.6-6.5	85	16.3	70	15.9	43	13.6	198	15.5
6.6-10.5	66	12.6	31	7.1	18	5.7	115	8.9
10.6-14.5	52	9.9	31	7.1	23	7.3	106	8.3
14.6-17.0	54	10.3	13	2.9	22	6.9	89	6.9
Non-adult	39	7.5	9	2.1	5	1.6	53	4.1
Total N	522		440		317		1279	

% of site total

Table 3. Number and percent overview of non-adult pathology by lesion.

Lesion	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%
CO&PH	522	65	12.5	440	41	9.3	317	31	9.8
THAL	522	3	0.6	440	0	0	317	0	0
LEH	522	83	15.9	440	43	9.8	317	13	4.1
EL&SPNBF	351	53	15.1	211	43	20.4	145	21	14.5
TB	522	4	0.8	440	2	0.5	317	1	0.3
METABOLIC	351	15	4.3	211	6	2.8	145	10	6.9
TRAUMA	522	7	1.3	440	5	1.1	317	5	1.6

% affected of observed by age group and site type; CO: cribra orbitalia, PH: porotic hyperostosis, THAL: thalassaemia, LEH: linear enamel hypoplasia, EL: endocranial lesions, SPNBF: sub-periosteal new bone formation, TB: tuberculosis.

Table 4. Number and percent of haematopoietic skeletal lesions (cribra orbitalia and porotic hyperostosis).

Age (yrs)	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%
Perinate	88	0	0	84	0	0	95	0	0
0.0-1.0	83	4	4.8	145	0	0	77	1	1.3
1.1-2.5	55	6	10.9	57	6	10.5	34	10	29.4
2.6-6.5	85	16	18.8	70	19	27.1	43	7	16.3
6.6-10.5	66	10	15.2	31	7	22.6	18	5	27.8
10.6-14.5	52	10	19.2	31	6	19.4	23	3	13.1
14.6-17.0	54	14	25.9	13	3	23.1	22	5	22.7
Non-adult	39	5	12.8	9	0	0	5	0	0
Total	522	65	12.5	440	41	9.3	317	31	9.8

% affected of observed by age group and site type

Table 5. Number and percent of haematopoietic skeletal lesions (cribra orbitalia and porotic hyperostosis).

Age (yrs)	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%	<i>Observed N</i>	<i>Affected n</i>	%
Perinate	88	0	0	84	0	0	95	0	0
0.0-1.0	83	0	0	145	0	0	77	0	0
1.1-2.5	55	6	10.9	57	7	12.3	34	0	0
2.6-6.5	85	19	22.4	70	12	17.1	43	1	2.3
6.6-10.5	66	18	27.3	31	7	22.6	18	5	27.8
10.6-14.5	52	22	42.3	31	13	41.9	23	4	17.4
14.6-17.0	54	17	31.5	13	4	30.8	22	2	9.1
Non-adult	39	1	2.6	9	0	0	5	1	20.0
Total	522	83	15.9	440	43	9.8	317	13	4.1

% affected of observed by age group and site type

Table 6. Number and percent of lesions indicative of inflammation/infection (sub-periosteal new bone formation, endocranial lesions).

Age (yrs)	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>
1.1-2.5	55	9	16.4	57	15	26.31	34	5	14.7
2.6-6.5	85	10	11.8	70	14	20	43	8	18.6
6.6-10.5	66	9	13.6	31	7	22.6	18	4	22.2
10.6-14.5	52	9	17.3	31	5	16.1	23	3	13.1
14.6-17.0	54	10	18.5	13	2	15.4	22	1	4.5
Non-adult	39	6	15.4	9			5		
Total	351	53	15.1	211	43	20.4	145	21	14.5

% affected of observed by age group and site type

Table 7. Number and percent of vitamin D and C deficiencies.

Age (yrs)	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>
Vitamin D deficiency									
1.1-2.5	55	4	7.3	57	2	3.5	34	2	5.9
2.6-6.5	85	1	1.2	70	1	1.4	43	0	0
6.6-10.5	66	1	1.5	31	1	3.2	18	0	0
10.6-14.5	52	0	0	31	0	0	23	0	0
14.6-17.0	54	1	1.9	13	0	0	22	0	0
Non-adult	39	1	2.6	9	0	0	5	0	0
Total	351	8	2.3	211	4	1.9	145	2	1.4
Vitamin C deficiency									
1.1-2.5	55	1	1.8	57	1	1.8	34	3	8.8
2.6-6.5	85	3	3.5	70	1	1.4	43	2	4.7
6.6-10.5	66	1	1.5	31	0	0	18	1	5.6
10.6-14.5	52	0	0	31	0	0	23	2	8.7
14.6-17.0	54	1	1.9	13	0	0	22	0	0
Non-adult	39	0	0	9	0	0	5	0	0
Total	351	6	1.7	211	2	0.9	145	8	5.6

% affected of observed by age group and site type

Table 8. Number and percent of trauma.

Age (yrs)	Major urban			Minor urban			Rural		
	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>	<i>Observed N</i>	<i>Affected n</i>	<i>%</i>
Perinate	88	0	0	84	0	0	95	0	0
0.0-1.0	83	1	1.2	145	0	0	77	1	1.3
1.1-2.5	55	1	1.8	57	1	1.8	34	0	0
2.6-6.5	85	0	0	70	0	0	43	1	2.3
6.6-10.5	66	0	0	31	0	0	18	0	0
10.6-14.5	52	2	3.8	31	3	9.7	23	1	4.3
14.6-17.0	54	2	3.7	13	1	7.7	22	2	9.1
Non-adult	39	0	2.3	9	0	0	5	0	0
Total	522	7	1.3	440	5	1.8	317	5	1.6

% affected of observed by age group and site type

Table 9. Fracture sites by skeletal element.

Element	Major urban		Minor urban		Rural		Total	
	<i>n</i>	<i>%</i>	<i>n</i>	<i>%</i>	<i>n</i>	<i>%</i>	<i>n</i>	<i>%</i>
Clavicle	1	14.3	1	20.0	3	60.0	5	29.4
Rib	1	14.3	0	0	0	0	1	5.9
Vertebra	1	14.3	0	0	0	0	1	5.9
Humerus	1	14.3	0	0	0	0	1	5.9
Radius	2	28.6	0	0	0	0	2	11.8
Ulna	0	0	0	0	1	20.0	1	5.9
Phalanx (hand)	0	0	1	20.0	0	0	1	5.9
Femur	0	0	1	20.0	0	0	1	5.9
Tibia	1	14.3	1	20.0	1	20.0	3	17.6
Phalanx (foot)	0	0	1	20.0	0	0	1	5.9
Total N	7		5		5		17	

% of total fractures by site




FIG. 1. Map of sites with non-adult remains included in survey.

MAJOR URBAN: 1. York; 2. Leicester; 3. Gloucester; 4. Cirencester; 5. Colchester; 6. London; 7. Winchester.

MINOR URBAN: 8. Ancaster; 9. Great Casterton; 10. Ashton; 11. Chesterton; 12. Baldock; 13. Dunstable; 14. Queenford Farm/Queensford Mill; 15. Springhead.

RURAL: 16. Bantymock Mine; 17. Watersmeet; 18. Huntsman's Quarry; 19. Babraham Institute; 20. Frocester; 21. Cannington; 22. Bradley Hill; 23. Catsgore; 24. Owselbury; 25. Dorchester By-pass; 26. Dewlish.




FIG. 2. Percentage distribution of ages-at-death.




FIG. 3. Overview of percentage distributions of pathological lesions (CO: cribra orbitalia, PH: porotic hyperostosis, EL: endocranial lesions, SPNBF: sub-periosteal new bone formation).




FIG. 4. Percentage distribution of haematopoietic lesions by age group.




FIG. 5. Percentage distribution of enamel hypoplasia by age group.




FIG. 6. Percentage distribution of sub-periosteal new bone formation and endocranial lesions by age group.




FIG. 7. Percentage distribution of vitamin D deficiency by age group.




FIG. 8. Percentage distribution of vitamin C deficiency by age group.




FIG. 9. Distribution of fracture locations by settlement type. Template adapted from Roksandic (2003, fig. 2).



FIG. 10. Digital radiograph of right clavicle with healed fracture (top) at the mid-shaft. Bottom is left clavicle for comparison. From Queensford Farm/Mill, skeleton 51/232. (*With permission from Oxfordshire Museums Service*)



FIG. 11. Right: left humerus with fracture at the mid-shaft, resulting in prominent angulation and changes to the proximal and distal epiphyseal ends. Left: right humerus for comparison. From Butt Road, Colchester, skeleton 595. (*With permission from Colchester and Ipswich Museums*)



FIG. 12. Infant left rib with fracture callus. From Winchester (Victoria Road East cemetery), skeleton 444. (*With permission from Hampshire Cultural Trust*)